



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Diminished nap effects on memory consolidation are seen under oral contraceptive use

Citation for published version:

Genzel, L, Bäurle, A, Potyka, A, Wehrle, R, Adamczyk, M, Friess, E, Steiger, A & Dresler, M 2014, 'Diminished nap effects on memory consolidation are seen under oral contraceptive use', *Neuropsychobiology*, vol. 70, no. 4, pp. 253-61. <https://doi.org/10.1159/000369022>

Digital Object Identifier (DOI):

[10.1159/000369022](https://doi.org/10.1159/000369022)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Neuropsychobiology

Publisher Rights Statement:

Authors' final peer reviewed manuscript as accepted for publication

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



**Diminished nap effects on memory consolidation are seen
under oral contraceptive use**

Lisa Genzel^{1#}, Anna Bäurle^{1*}, Alina Potyka^{1*}, Renate Wehrle¹, Marek Adamczyk¹,
Elisabeth Friess¹ D, Axel Steiger¹, Martin Dresler^{1,2}

¹Max Planck Institute of Psychiatry, Munich/Germany
²Donders Institute for Brain, Cognition and Behaviour, Radboud University Medical
Center, Nijmegen/Netherlands

*The authors contributed equally to the study.

The work was performed at the Max Planck Institute of Psychiatry Munich/Germany

Correspondence: Dr. med Lisa Genzel, Max Planck Institute of Psychiatry,
Kraepelinstr. 2-10, 80804 Munich, Germany; genzel@mpipsykl.mpg.de; Tel: +49 89 306 22
386 Fax: +49 89 306 22 552

Current Address: Dr. med Lisa Genzel, Centre for Cognitive and Neural Systems,
University of Edinburgh, 1 George Square, Edinburgh EH8 9JZ, UK; LGenzel@ed.ac.uk;
Tel: +44 131 650 4571 Fax: +44 131 651 1835

Abstract

Many young females take exogenous hormones as oral contraceptive (OC); a condition rarely controlled for in studies on sleep and memory consolidation even though sex hormones influence consolidation. This study investigated the effects of OCs on sleep-related consolidation of a motor and declarative task, utilizing a daytime nap protocol.

Fifteen healthy, young females taking OCs came to the sleep lab for three different conditions: nap with previous learning, wake with previous learning and nap without learning. They underwent each condition twice, once during the 'pill active' weeks and once during the 'pill free week', resulting in six visits.

In all conditions, participants showed a significant offline consolidation effect, independent of pill week or nap/wake condition. There were no significant differences in sleep stage duration, spindle activity or spectral EEG frequency bands between naps with or without learning condition.

The present data showed a significant offline enhancement in memory irrespective of potential beneficial effects of a nap. In comparison to previous studies this may suggest that the use of oral contraceptives may enhance offline memory consolidation in motor and verbal tasks per se. These results stress the importance to control for the use of OCs in studies focusing on memory performance.

Keywords: nap, sleep, oral contraceptives, procedural motor learning, declarative verbal memory, estrogen, progesterone, pill, spindles

1 **Introduction**

2 The evidence for a beneficial role of sleep on memory consolidation is becoming
3 stronger [1;2]. However, several studies show diverging results including a lack of
4 improvement following sleep [3-8]. A possible explanation for this confusion may be a
5 disregard of additional confounding factors. For one, studies rarely control for sex, menstrual
6 cycle or the use of oral contraceptives (OCs). The hormones estrogen and progesterone have a
7 wide range of effects on sleep as well as on memory. On the molecular and synaptic level,
8 estrogen positively influences the hippocampus and other memory related brain areas, by
9 inducing a beneficial environment for memory encoding and consolidation [9;10]. On the
10 behavioral level, it is important to distinguish between tasks in which males typically show an
11 advantage (e.g. spatial) and tasks in which females typically show an advantage (e.g. verbal,
12 fine motor) [11]. This has clearly been demonstrated in animal models; however results in
13 human studies are more variable. These latter tasks are positively influenced by the hormones
14 estrogen and progesterone. In contrast, tasks with a male advantage are affected negatively by
15 the same hormones [12-16]. Furthermore, use of OCs influences memory encoding: Females
16 showed enhanced verbal memory during the active OC phase [17]. Another study presented
17 that after sleep deprivation, females in the follicular phase performed worse on different
18 cognitive tests than females in the luteal phase or taking OCs [18]. Wharton and colleagues
19 [19] could show by comparing different OC products that the androgenic activity in OCs
20 influence mental rotation task performance, a typical “male” task. Not only memory, but also
21 sleep is influenced by exogenous hormones. Females taking OCs have less slow wave sleep,
22 increased stage 2 sleep, shorter REM onset latency and more REM sleep than naturally
23 cycling women [20;21].

24 In a previous study we could demonstrate a sex and menstrual cycle effect on sleep-
25 related memory consolidation of “female” tasks [22]. While male subjects benefitted from a

nap in verbal and in motor learning, females did so only during the mid-luteal phase with high levels of estrogen and progesterone, however not during the early-follicular phase with low levels of the respective hormones. Effects in motor learning were correlated with hormonal levels of progesterone, and effects in verbal learning with levels of estrogen in the participants. Sleep spindles showed a similar effect. Spindle activity increased upon learning in males, whereas in females it increased only during the mid-luteal phase, matching the learning behavior. Furthermore, sleep spindle density and frequency correlated with estrogen [22].

The majority of studies on human memory functions investigate healthy young subjects. At the same time many young females take OCs – around 72% of all 18-29 year old females in Germany [23] – therefore are under the influence of exogenous estrogen and progesterone. However, this condition is rarely controlled for or even regarded as potential confound.

Aim and hypothesis

To investigate if OC use in participants of sleep and memory studies may confound the outcome of these studies, we investigated the effects of OCs on verbal and motor memory consolidation during a daytime nap. A nap has been shown to be as effective for memory consolidation as a whole night of sleep for these tasks, but has the advantage of avoiding time of day or stress via sleep deprivation as confounds [24-34]. Females taking OCs underwent three conditions – a nap with learning, wake with learning, and nap without learning. Participants did so once during a pill week and once during the regularly recurring pill free week, resulting in six visits altogether. Based on our previous finding of strongest enhanced memory consolidation during the third week of the natural menstrual cycle, with highest levels of estrogen and progesterone, we expected to see a similar strong effect in the participants when taking OCs. We hypothesized that participants taking OCs would show enhanced memory consolidation in both tasks.

Materials and Methods

Participants

The participants were healthy female volunteers (n=15) aged 18-30 years taking oral contraceptives. They were recruited mainly via the local medical school and were paid for the participation in the study. All participants were first screened for psychiatric, physical, or sleep disorders with a semi-structured interview, physical examination and the Pittsburgh Sleep Quality Index [35]. Additionally, we obtained urinary-drug-screening and routine blood tests (blood cell count, electrolytes, liver and kidney function, thyroid hormones). Further exclusion criteria were: shift-work at night, a transmeridian flight, any medical treatment during the last three months, substance abuse (assessed via oral question and urinary drug screening), professional piano playing (more than 5 years intensive training), professional type writing, extreme chronotypes (scores of >70 and <30, assessed via the D-MEQ [36;37]) and regular naps (>2 nap/month). Professional pianists and typists were excluded since their baseline tapping performance would be significantly higher than the other subjects, which could affect offline improvement. All participants took one of two types of oral contraceptives (OCs): Valette (Bayer Austria GmbH Vienna/Austria) and Belara (Gruenthal GmbH Aachen/Germany). Both OCs had equal amount of estradiol (0.03 mg ethinylestradiol), and different gestagens (2 mg dienogest (Valette) and 2mg chlormadinoacetat (Belara) respectively). OC intake was for at least one year prior to the study, following the generally recommended scheme of three weeks of daily intake followed by a 'discontinuation' week during which menses may occur.

The participants agreed to have regular sleep patterns throughout the experiment and kept a sleep diary for each week preceding a study block. The Ethics Committee of the Ludwig Maximilian University, Faculty of Medicine, Munich/Germany, approved the research project. The experiments were undertaken with the understanding and written

consent of each subject, and the study conforms to The Code of Ethics of the World Medical Association.

Procedures

All subjects underwent 6 study days (see Figure 1): nap with learning, wake with learning, nap without learning; each of the three conditions once during the ‘pill active’ phase during OC intake (second week of the pill cycle) and once during the monthly recurring ‘pill-free’ week. The order of all 6 conditions was randomized and balanced between participants. The conditions were separated by $27d \pm 18$ with a range of 10-86 days. In addition, in each experimental condition blood for hormonal analysis was drawn from a peripheral vein after the subjects arrived at the lab.

The nap-protocol used in this study, has been established and used previously [22;24;34]. During study days the subjects arrived at 13.00 h; they first completed the D2 Concentration test (D2) [38] and the Stanford Sleepiness Scale (SSS) [39] followed by the learning phase of a verbal paired associate task [32] and a sequential finger tapping task [40]. During the learning phase and the retest we conducted the four learning and alertness tasks in a randomized order to avoid a confounding effect of a reciprocal interaction between the tasks [41]. We had three “stations” (SSS and D2, tapping, word pairs), which resulted in 6 different orders. The subjects were pseudo randomly (so randomly but then balancing across participants with 2-3 participants having the same order) assigned one of the 6 order sequences. Subsequently at around 14.00h the subjects were informed to which condition they had been assigned: participants in the WAKE-condition (L-Wake) watched a non-emotionally-arousing movie until retest; in participants in the NAP-condition the electrodes were placed, the lights were turned off, and the subjects were allowed to sleep for

approximately 60 min (L-NAP). A 60 min nap duration was chosen matching previous studies [22;24;34] so that most subjects would have naps containing stage 2 sleep and slow wave sleep. At around 16:30 h or at least 30 min after awakening from the approx. 1-hour-nap, all subjects completed the D2 test, the SSS, and the retest, after which they returned home.

During the nap without learning condition the participants arrived at 14:00, filled out the D2 and SSS and took a nap at the same time of day (C-NAP), with polysomnographic recordings but without learning tasks. This condition consisted solely of a nap without learning to investigate changes in sleep induced by learning (C-NAP vs. L-Nap).

The participants were instructed to refrain from rehearsal of the tasks and to keep a regular sleep cycle throughout the weeks of the experiment. In addition the participants kept a sleep diary for a week preceding each study block. During this week they went to bed between 23.00h and 1.00h and woke between 7.00h and 9.00h; during the three nights prior the study day the bedtime changed to 23.00 – 24.00h and the wake-time to 7.00 – 8.00h.

Figure 1 please near here

Hormone measures

Directly after the participants arrived at the sleep lab blood was drawn for hormonal analysis. Immediately after the draw the test tubes (serum-tubes with clot activator, 7.5 ml, from Sarstedt Nuembrecht/Germany 01.1601.001) were centrifuged and transferred to the in-house lab for analysis, or refrigerated (~4°C) until analysis could be performed. Hormones – 17 beta estradiol and progesterone – were measured by electrochemiluminescence, with an Elecsys 2010 analyzer (Roche Diagnostics, Basel/Switzerland). Functional sensitivity for 17 beta estradiol was 12 pg/mL, and for progesterone 0.15 ng/mL. In our lab it was only possible

1 to measure the levels of endogenous hormones estrogen and progesterone and not of the
2 exogenous OC hormones. Reported pharmacological properties and measurements for Valetta
3 are for dienogest: maximum plasma concentration $51.6 \pm 9.5 \text{ ng/ml}$ reached in $2.4 \pm 1.4 \text{ h}$, steady
4 state after daily intake 1.5 fold serum levels, 96% bioavailability, 10% plasma free form, 90%
5 bound to albumin, $9.3 \pm 1.8 \text{ h}$ half-life, $3.66 \pm 0.71 \text{ L/h}$ clearance; and for ethinyloestradiol:
6 maximum plasma concentration reached in 1.5-4h steady state after daily intake 2 fold serum
7 levels, 44% bioavailability, 98.5% bound to albumin, $11.7 \pm 6.5 \text{ h}$ half-life, 5 mL/min/kg
8 metabolic clearance.

9 10 *Polysomnographic recording parameters*

11 Polysomnographic data were recorded in all nap conditions; stored and analyzed with a
12 digital recorder (Comlab 32 Digital Sleep Lab, Brainlab V 3.3 Software, Schwarzer GmbH,
13 Munich, Germany). We recorded scalp EEG from the C3 and C4 derivations referenced
14 against the contralateral mastoid (filtered from 0.5 to 70 Hz), and further electrooculograms
15 (EOG) and mental/submental electromyogram (EMG), with a sampling rate of 250 Hz.

16 17 *Learning Tasks*

18 All subjects learned two tasks; one declarative (verbal) and one procedural task (motor).

19 The tool for declarative memory analysis was a paired associates learning task. We used
20 paralleled standardized word lists consisting of 40 related word pairs (e.g. Nanny – Stroller),
21 with additional 2 dummy pairs in the beginning and at the end to avoid inclusion of primacy
22 and recency effects [32]. In the learning condition the word pairs were first presented for 5 s
23 each, and immediately after the list presentation a cued recall followed in which the
24 participant was asked to type each matching noun after being shown the first word of the pair.
25 If the participant was not able to recall the right word, the correct answer was displayed. Thus

every participant saw the correct pairing twice, once in the learning phase and once during retest. This method aims to avoid differences in exposure to the learning material by differences in recall performance. Each word pair was cued once. In the retest condition (delayed recall after approx. 3.5hrs following nap / wake condition), the cue words were given once and the number of correctly known word pairs was obtained by the experimenter to compensate for spelling errors. At the training and retest condition the subject had unlimited time to respond to the cued recall. In order to measure sleep-related consolidation we used absolute change in performance from learning to retest (e.g. performance at learning 15 correct word pairs and performance at retest 20 correct word pairs resulted in consolidation measure of 5).

To test procedural motor memory, we employed a sequential finger tapping task [40]. This task required participants to press four numeric keys on an altered computer keyboard with their non-dominant hand, repeating the five element sequence (e.g. 4-1-3-2-4) as quickly and accurately as possible for a period of 30 s. Four different sequences were used in the experiments. To exclude any working memory component on the task, the numeric sequence was displayed on the screen. For every trial the computer noted the number of complete sequences achieved, the number of errors made, and the number of correct sequences typed. The learning phase consisted of twelve trials of 30 s interrupted by 20 s rest periods, while at retest the subjects had to complete four trials. As score we used the number of the correctly tapped sequences during the period of 30 s, which incorporates the accuracy and speed performance. End-training performance consisted of the average score from the last three trials of the training, while retest performance was composed of the average score from all four retest trials. To measure sleep-related consolidation, end-training performance was used as baseline and the change to retest performance was divided by the end-training performance

(e.g. performance at learning 20 correctly typed sequences per 30s and performance at retest 25 correctly typed sequences per 30s resulted in consolidation measure of 25%).

Sleep data analysis

For sleep data analysis, independent professionals scored the sleep stages using standard criteria [42]. The scorers were blind of the study design. Additionally, the EEG of the experimental naps (L-Nap, C-Nap), contra-lateral to the typing hand, underwent a spectral analysis through a fast-fourier-transformation using in-house software. The EEG was digitally filtered from 0.53 to 30 Hz (24dB/octave) after sweeps with visually identified EEG artifacts had been carefully removed. Power spectra were derived from 2 s windows, shifted for 1 s and averaged per epoch of 30 s. Frequency bands (based on summed power values) were calculated for the delta (0.53 - 4 Hz), theta (4.5 – 8 Hz), alpha (8.5 – 12 Hz), sigma (12.5 – 16 Hz), and beta (16.5 – 20 Hz) frequency range.

Sleep spindle analysis

An automated algorithm detected the sleep spindles. The algorithm first removes periods of EEG signal with muscle artifacts and strong alpha frequencies. Afterwards an individual spindle threshold is set for each channel and spindles are identified with continuous wavelet transformation. For a more detailed description of the analysis see supplementary materials. Analyzed parameter was spindle activity (SpA; mean spindle amplitude×mean spindle duration). We used SpA since it well reflects the intensity of the spindle process [43;43;44;44;45;45].

Statistical Analysis

For statistical analysis of offline memory consolidation, each an ANOVA was performed for the verbal and motor consolidation measures with the within-subject factors week (OC/OC-free) and condition (nap/wake). In addition, the change in performance from the end of the learning phase to retest after sleep or wake for both tasks was tested via paired T-tests considering a bonferroni corrected statistical threshold ($p < 0.05/4$). For the polysomnographic data, we performed each a MANOVA with repeated measures of (a) the duration of sleep stages, (b) the EEG frequency bands and (c) spindle activity, with within-subjects factors “naps” (factor levels L-NAP and C-NAP) and week (OC/OC-free). The alertness data (D2, SSS) and the absolute end-training performance for both learning tasks were analyzed with each a MANOVA with within-subjects factors week (OC/OC-free) and condition (nap/wake). The hormone values of progesterone and estrogen were correlated with overnight change in memory performance. Alpha was set at 0.05.

Results

There was no week or condition effect on the absolute end-training performance of both the declarative and the motor task (declarative: condition (L-Nap vs. L-Wake): $F_{1,14}=.13$, $P>.7$; week (OC/OC-free): $F_{1,14}=.001$, $P>.9$; condition*week: $F_{1,14}=.11$, $P>.7$; motor: condition (L-Nap vs. L-Wake): $F_{1,14}=.44$, $P>.8$; week (OC/OC-free): $F_{1,14}=.08$, $P>.7$; condition*week: $F_{1,14}=2.39$, $P>.1$), demonstrating that all subjects started from comparable baseline levels. There was a practice effect (baseline/retest) on the concentration task but not on the sleepiness scale: MANOVA with factors test (baseline/retest), week (OC/OC-free), condition (L-Nap/L-Wake), and their interactions showed a significant effect for test (all $F_{2,12}=8.74$, $P=0.005$, D2: $F_{1,13}=14.55$, $P=0.002$, SSS: $F_{1,13}=2.02$, $P>0.1$) but no interaction or factor effects for week and condition (all $P>0.05$). An ANOVA with the factors test (baseline/retest), week (OC/OC-free), condition (L-Nap/L-Wake), and their interactions showed a significant effect for test (verbal learning: $F_{1,14}=69.019$, $P<0.001$, motor learning: $F_{1,14}=43.404$, $P<0.001$) but no interaction or factor effects for week and condition (all $P>0.05$). For both tasks a significant increase from end-training performance to post nap/wake retest performance was seen in all 4 conditions (all $P<.008$ with corrected threshold at $P<.0125$, see Table 1). The ANOVAs for motor learning (condition (L-Nap/ L-Wake): $F_{1,14}=.031$, $P>.8$; week (OC/OC-free): $F_{1,14}=2.282$, $P>.15$; condition*week: $F_{1,14}=.355$, $P>.5$) as well as verbal learning (condition (L-Nap/L-Wake): $F_{1,14}=.016$, $P>.9$; week (OC/OC-free): $F_{1,14}=.225$, $P>.6$; condition*week: $F_{1,14}=1.377$, $P>.2$) showed no significant differences in the offline consolidation measures between any of the different conditions (see Figure 2). This remained the same if relative instead of absolute increase was used for verbal learning (condition (L-Nap/L-Wake): $F_{1,14}=.363$, $P>.5$; week (OC/OC-free): $F_{1,14}=.090$, $P>.7$; condition*week: $F_{1,14}=.322$, $P>.5$). There was no significant offline change in errors in the motor task or a condition/week effect on errors indicating that the increase in general motor performance was

1 due to an increase in speed (ANOVA with factors test (baseline/retest), week (OC/OC-free),
2 condition (L-Nap/L-Wake), and their interactions showed no interaction or factor effects for
3 test, week and condition (all $P > 0.45$).

4
5 *Figure 2 and Table 1 please near here*

6
7 All subjects fell asleep during their naps with an average sleep duration of > 60 min
8 (with average light-out of ~ 90 min). Polysomnographic data revealed allocation of sleep
9 stages with mainly stage 2 sleep and SWS, and additionally a small amount of REM sleep in
10 some subjects. There was no effect of conditions (C-Nap/L-Nap/OC/OC-free) on sleep stage
11 distribution or data from spectral analysis of the sleep EEG (see Table 2). There were no
12 condition or OC phase effects on sleepiness and concentration at the learning phase or at
13 retest (condition (L-Nap vs. L-Wake): $F_{4,10} = .899$, $P > .5$; week (OC/OC-free.): $F_{4,10} = 1.566$,
14 $P > .2$; condition*week: $F_{4,10} = .552$, $P > .7$). No significant effect of week or condition on spindle
15 activity could be found (condition: $F_{1,14} = .019$, $P > .8$; week: $F_{1,14} = .227$, $P > .6$; condition*week:
16 $F_{1,14} = .028$, $P > .8$); this remained true for other spindle measure in sleep stage 2 as well as
17 considering all NREM (see suppl. materials). Endogenous hormonal levels of all six
18 conditions are presented in Table 3.

19
20 *Table 2 and 3 please near here*

21
22 The change in tapping performance and word pairs did not correlate significantly with
23 the amount of each sleep stage (stage 2, SWS, REM, TST) or with sleep spindle activity
24 during the naps (all 2-tailed, $r < .3$, $P > .15$). We did find a significant positive correlation

1 between change in word pairs and endogenous estrogen across all conditions (1-tailed, $r=.358$,
2 $P<.003$). However, the correlation seemed dominated by one outlier. After exclusion of the
3 outlier the correlation still was significant, but only 1-tailed (1-tailed, $r=.235$, $P<.05$). The
4 change in word pairs did not correlate with progesterone and the change in tapping did not
5 correlate with any of the hormone values (all 2-tailed, $r<.14$, $P>.25$).

6 Sample size and power calculation are presented in the supplementary materials.

Discussion

This study investigated the effects of oral contraceptives (OC) on offline memory consolidation (= all consolidation processes, which occur when the person is not actively engaged in learning). Participants taking a contraceptive pill performed at a significantly higher level during retest four hours after the learning session. This improvement occurred regardless of an interim nap of roughly 60 minutes or staying awake in the same period. This finding also occurred irrespective of OC week (active OC uptake or (monthly) OC-free week).

In a previous study utilizing the same tasks and procedures, we had investigated the effects of the menstrual cycle on memory consolidation [22]. In the menstrual cycle study the participants started at a similar behavioral baseline as in the present study, however only the females in the nap condition during the mid-luteal phase (high with estrogen) managed to increase their performance by 7 word pairs, while all other groups/conditions (men or females in the follicular phase) only knew roughly 4 word pairs more during the retest (for visual comparison see supplementary figure 1). This might indicate that the increase of 8 word pairs in the current study – regardless of OC phase or nap/wake condition – may represent a comparable strong improvement, possibly connected to the exogenous and endogenous hormonal levels. As seen in the previous study [22], we again found a correlation between endogenous estrogen and change in word pairs. Regrettably only endogenous and not OC hormone levels could be measured in our lab, since most likely the strong improvement was induced by the endogenous as well as the exogenous hormones.

Independent of the length or content of the word lists used, sleep related effects on verbal memory usually seem to occur in a similar range. Lists with 40 word pairs (based on [32] as used here) are the most common tool in studies investigating effect of sleep on declarative memory. Irrespective of the length of sleep (nap or whole night condition), the

1 offline change reported is usually in the range of -2 to +5 word pairs [24-32]. Only the
2 studies by Tucker and colleagues [33;34] reported a higher offline change of around 8 word
3 pairs as was similarly found in the present data on OC use, as well as during the luteal phase
4 in women [22]. However, it would be beneficial to replicate this study with a whole night of
5 sleep to confirm that the length of sleep does not influence offline change.

6 A similar effect is seen in the tapping performance. On average, participants increase
7 their performance by roughly 0-5% after wake and 10-30% after sleep [22;27-31;40;46-55].
8 Regardless of OC phase or nap/wake condition, the increase reported here was 10 to 17%
9 similar to previous data seen only after sleep.

10 A positive effect of OCs on memory encoding has been shown previously. Participants
11 taking OCs performed better at a verbal task during immediate testing – not delayed as in this
12 study – than natural cycle women [17;17]. In this study we did not find an effect of OC-phase
13 (active OC intake or OC-free interval) on memory. While some studies do report a OC-phase
14 effect [17;56], other studies do not find such an effect [57-59]. It does not seem too surprising
15 that there was no phase effect on memory, if one considers the range of absolute hormone
16 values. While our subjects did show a significant rebound-effect in estrogen during the OC-
17 free week, the values of endogenous estrogen consistently remained low in comparison to
18 women with normal menstrual cycles (ranges: OC 9-50 pg/ml, menstrual cycle 55-155 pg/ml
19 see [22]) while exogenous estrogen levels were most likely high.

20 There seemed to be no additional benefit of a nap on memory consolidation in this study.
21 There are different possible explanations for this finding. One likely assumption is that the
22 hormones in the OCs boost the consolidation in such a way that no additional benefit of sleep
23 was possible. Another possibility is that a ceiling effect was reached in the tasks themselves.
24 Further, it is also possible that estrogen increased plasticity during encoding and that
25 increased encoding masked or influenced the effects of sleep on consolidation. Especially

1 since it has been reported previously that pre-sleep performance levels can influence sleep
2 related benefit [60] .

3 A wide range of effects of estrogen and progesterone on the hippocampus and other
4 brain areas important for memory has been observed. An estrogen influence on plasticity was
5 evidenced after exogenous estradiol administration in ovariectomised rats by increases in
6 neurogenesis [61], neural network connectivity and synaptic transmission [9]. Furthermore,
7 estrogen increases glucose transport, glycolysis and mitochondrial function to provide the
8 ATP necessary for energetic demand as seen in non-human primates and after exogenous
9 estradiol administration in ovariectomised rats [9]. Estrogen affects cell morphology, synapse
10 formation, signaling and excitability in the hippocampal formation [62-64]. In the
11 hippocampus and the medial prefrontal cortex estrogen increases dendritic spines, and an
12 increase in spine density has been associated with learning and memory [9;64]. Estrogens
13 upregulate adult hippocampal neurogenesis and synaptic protein levels in the hippocampus as
14 well as enhance synaptic NMDA receptor current and the magnitude of long-term potentiation,
15 a cellular correlate of learning and memory [14-16].

16 It seems that in humans as well as rodents estrogen affects different types of memory
17 differently [11;65;66]. In general, memory can be divided into tasks in which females show
18 an advantage (fine motor, verbal, object location etc.) as well as tasks in which males show an
19 advantage (mainly spatial) [67-70]. “Female” tasks seem to be positively influenced by the
20 hormones estrogen and progesterone, while “male” tasks seem to be negatively influenced
21 [12-16]. In both types of tasks a menstrual waxing- and waning effect can be seen. On tasks in
22 which women typically score better than men, women perform better during mid-luteal phase
23 (high estrogen and progesterone) than within menstrual phase (low estrogen and
24 progesterone). On tasks in which men typically outperform women, women do best during
25 menses [13;71;72].

Caveats

It is important to note that this study does not intend to advertise OCs as neuroenhancers. For one, we did not perform a placebo controlled, double blind cross-over study, which would be needed to be able to attempt this conclusion. Secondly, our sample size may also have been too small to detect more subtle effects, however we did not even see a trend in the data and the sample size is comparable with most studies investigating sleep related consolidation. Thirdly, we did not investigate the effect of OC use on “male” learning tasks. Since female hormones actually exhibit negative effects on memory tasks in which males outperform females, the offline consolidation of those tasks may actually be reduced by OC use. Instead this study attempts to underline the importance to acknowledge OC use as an influencing factor in sleep and memory research, which should be controlled or manipulated. A further caveat is that we did not perform an adaption nap, which could have influenced the result.

Conclusion

We could show that female participants taking OCs experienced a significant and rather large improvement during offline consolidation in a verbal and a fine motor task independent of nap/wake condition. It is tempting to speculate that this already strong enhancement in comparison to other studies was caused by the OCs and masked any potential sleep effects. These results are important pilot findings and should be confirmed with a placebo controlled, double blind cross-over study. But they do point towards the importance to control for OC use in studies investigating memory effects. Such effects may also hold responsible for some of the discrepancies in previously published results.

Acknowledgments

1 We thank our participants and the sleep lab team for expert data handling.

2

Reference List

- 1 Diekelmann S, Born J: The memory function of sleep. *Nat Rev Neurosci* 2010;11:114-126.
- 2
- 3 2 Rattenborg NC, Martinez-Gonzalez D, Roth II TC, Pravosudov VV:
- 4 Hippocampal memory consolidation during sleep: a comparison of mammals and
- 5 birds. *Biological Reviews* 2011;86:658-691.
- 6
- 7
- 8 3 Cai D, Rickard T: Reconsidering the role of sleep for motor memory. *Behav*
- 9 *Neurosci* 2009;123:1153-1157.
- 10
- 11 4 Keisler A, Ashe J, Willingham DT: Time of day accounts for overnight
- 12 improvement in sequence learning. *Learn Mem* 2007;14:669-672.
- 13
- 14 5 Nemeth D, Janacek K, Londe Z, Ullman M, Howard D, Howard J: Sleep has
- 15 no critical role in implicit motor sequence learning in young and old adults.
- 16 *Experimental Brain Research* 2010;201:351-358.
- 17
- 18 6 Rickard TC, Cai DJ, Rieth CA, Jones J, Ard MC: Sleep does not enhance
- 19 motor sequence learning. *Journal of Experimental Psychology: Learning, Memory,*
- 20 *and Cognition* 2008;34:834-842.
- 21
- 22 7 Song S, Howard Jr JH, Howard DV: Sleep does not benefit probalistic motor
- 23 sequence learning. *J Neurosci* 2007;27:12475-12483.
- 24
- 25 8 Conte F, Ficca G: Caveats on psychological models of sleep and memory: A
- 26 compass in an overgrown scenario. *Sleep Med Rev*.
- 27
- 28 9 Brinton RD: Estrogen-induced plasticity from cells to circuits: predictions for
- 29 cognitive function. *Trends in Pharmacological Sciences* 2009;30:212-222.
- 30
- 31 10 Mizuno K, Giese KP: Towards a molecular understanding of sex differences in
- 32 memory formation. *Trends in Neurosciences* 2010;33:285-291.
- 33
- 34 11 Andreano JM, Cahill L: Sex influences on the neurobiology of learning and
- 35 memory. *Learn Mem* 2009;16:248-266.
- 36
- 37 12 Miles C, Green R, Sanders G, Hines M: Estrogen and memory in a transsexual
- population. *Horm Behav* 1998;34:199-208.
- 13 Maki PM, Rich JB, Shayna Rosenbaum R: Implicit memory varies across the
- menstrual cycle: estrogen effects in young women. *Neuropsychologia* 2002;40:518-
- 529.
- 14 Barha CK, Galea LAM: Influence of different estrogens on neuroplasticity and
- cognition in the hippocampus. *Biochimica et Biophysica Acta* 2010;1056-1067.
- 15 Mukai H, Kimoto T, Hojo Y, Kawato S, Murakami G, Higo S, Hatanaka Y,
- Ogiue-Ikeda M: Modulation of synaptic plasticity by brain estrogen in the
- hippocampus. *Biochimica et Biophysica Acta* 2010;1030-1044.

- 16 Smith CC, Vedder LC, McMahon LL: Estradiol and the relationship between dendritic spines, NR2B containing NMDA receptors, and the magnitude of long-term potentiation at hippocampal CA3-CA1 synapses. *Psychoneuroendocrinology* 2009;34S:S130-S142.
- 17 Mordecai KL, Rubin LH, Maki PM: Effects of menstrual cycle phase and oral contraceptive use on verbal memory. *Horm Behav* 2008;54:286-293.
- 18 Wright KP Jr, Badia P: Effects of menstrual cycle phase and oral contraceptives on alertness, cognitive performance, and circadian rhythms during sleep deprivation. *Behav Brain Res* 1999;103:185-194.
- 19 Wharton W, Hirshman E, Merritt P, Doyle L, Paris S, Gleason C: Oral contraceptives and androgenicity: Influences on visuospatial task performance in younger individuals. *Experimental and Clinical Psychopharmacology* 2008;16:156-164.
- 20 Baker FC, Mitchell D, Driver HS: Oral contraceptives alter sleep and raise body temperature in young women. *Eur J Physiol* 2001;424:729-737.
- 21 Baker FC, Driver HS: Circadian rhythms, sleep, and menstrual cycle. *Sleep Med* 2007;8:613-622.
- 22 Genzel L, Kiefer T, Renner L, Wehrle R, Kluge M, Grözinger M, Steiger A, Dresler M: Sex and modulatory menstrual cycle effects on sleep related memory consolidation. *Psychoneuroendocrinology* 2012;37:987-989.
- 23 TNS Emnid for BZgA: Verhuetungsverhalten Erwachsener; 2012.
- 24 Backhaus J, Junghanns K: Daytime naps improve procedural motor memory. *Sleep Med* 2006;7:508-512.
- 25 Genzel L, Dresler M, Wehrle R, Grözinger M, Steiger A: Slow wave sleep and REM sleep awakenings do not affect sleep dependent memory consolidation. *Sleep* 2009;32:302-310.
- 26 Gorfine T, Yeshurun Y, Zisapel N: Nap and melatonin-induced changes in hippocampal activation and their role in verbal memory consolidation. *J Pineal Res* 2007;43:336-342.
- 27 Marshall L, Helgadóttir H, Mölle M, Born J: Boosting slow oscillations during sleep potentiates memory. *Nature* 2006;444:610-613.
- 28 Marshall L, Kirov R, Brade J, Mölle M, Born J: Transcranial electrical currents to probe EEG brain rhythms and memory consolidation during sleep in humans. *PLoS ONE* 2011;6:e16905.
- 29 Rasch B, Pommer J, Diekelmann S, Born J: Pharmacological REM sleep suppression paradoxically improves rather than impairs skill memory. *Nat Neurosci* 2009;12:396-397.

- 1 30 Wilhelm I, Diekelmann S, Molzow I, Ayoub A, Mölle M, Born J: Sleep
2 selectively enhances memory expected to be of future relevance. The Journal of
3 Neuroscience 2011;31:1563-1569.
- 4 31 Moroni F, Nobili L, Curcio G, De Carli F, Tempesta D, Marzano C, De
5 Gennaro L, Mai R, Francione S, Lo Russo G, Ferrara M: Procedural learning and
6 sleep hippocampal low frequencies in humans. NeuroImage 2008;42:911-918.
- 7 32 Plihal W, Born J: Effects of early and late nocturnal sleep on declarative and
8 procedural memory. J Cogn Neurosci 1997;9:534-547.
- 9 33 Tucker M, Fishbein W: The impact of sleep duration and subject intelligence
10 on declarative and motor memory performance: how much is enough? J Sleep Res
11 2009;18:304-312.
- 12 34 Tucker MA, Hirota Y, Wamsley EJ, Lau H, Chaklader A, Fishbein W: A
13 daytime nap containing solely non-REM sleep enhances declarative but not
14 procedural memory. Neurobiol Learn Mem 2006;86:241-247.
- 15 35 Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ: The Pittsburgh
16 sleep quality index: A new instrument for psychiatric practice and research.
17 Psychiatry Res 1989;28:193-213.
- 18 36 Horne JA, Osterberg O: A self-assessment questionnaire to determine
19 morningness-eveningness in human circadian rhythms. Int J Chronobiol 1976;4:97-
20 110.
- 21 37 Griefahn B, Kuenemund C, Broede P, Mehnert P: Zur Validitaet der deutschen
22 Uebersetzung des Morningness-Eveningness-Questionnaires von Horne und
23 Oestberg. Somnologie 2001;5:71-80.
- 24 38 Brickenkamp R: Test d2, Aufmerksamkeits-Belastungs-Test. ed 9, Göttingen,
25 Bern, Toronto, Seattle, Hogrefe, 2002.
- 26 39 Hoddes E, Zarcone V, Smythe H, Phillips R, Dement W: Quantification of
27 sleepiness: a new approach. Psychophysiology 1973;10:431-436.
- 28 40 Walker MP, Brakefield T, Morgan A, Hobson JA, Stickgold R: Practice with
29 sleep makes perfect: sleep-dependent motor skill learning. Neuron 2002;35:205-211.
- 30 41 Brown RM, Robertson EM: Off-Line Processing: Reciprocal Interactions
31 between Declarative and Procedural Memories. J Neurosci 2007;27:10468-10475.
- 32 42 Rechtschaffen A, Kales A: A manual of standardized terminology, techniques
33 and scoring system for sleep stages of human subjects. ed 3, Bethesda, MD, U.S.
34 Department of Health, Education and Welfare, Public Health Services, 1968.
- 35 43 Schabus M, Gruber G, Parapatics S, Sauter C, Klösch G, Anderer P, Klimesch
36 W, Saletu B, Zeitlhofer J: Sleep spindles and their significance for declarative
37 memory consolidation. Sleep 2004;27:1479-1485.

- 1 44 Schabus M, Hödlmoser K, Gruber G, Sauter C, Anderer P, Klösch G,
2 Parapatics S, Saletu B, Klimesch W: Sleep spindle-related activity in the human
3 EEG and its relation to general cognitive and learning abilities. *Eur J Neurosci*
4 2006;23:1738-1746.
- 5 45 Schabus M, Hödlmoser K, Pecherstorfer T, Anderer P, Gruber G, Parapatics S,
6 Sauter C, Klösch G, Klimesch W, Saletu B, Zeitlhofer J: Interindividual sleep
7 spindle differences and their relation to learning-related enhancements. *Brain Res*
8 2008;1191:127-135.
- 9 46 Fischer S, Hallschmid M, Elsner AL, Born J: Sleep forms memory for finger
10 skills. *Proc Natl Acad Sci* 2002;99:11987-11991.
- 11 47 Doyon J, Korman M, Morin A, Dostie V, Tahar A, Benali H, Karni A,
12 Ungerleider L, Carrier J: Contribution of night and day sleep vs. simple passage of
13 time to the consolidation of motor sequence and visuomotor adaptation learning.
14 *Experimental Brain Research* 2009;195:15-26.
- 15 48 Korman M, Doyon J, Doljansky J, Carrier J, Dagan Y, Karni A: Daytime sleep
16 condenses the time course of motor memory consolidation. *Nat Neurosci*
17 2007;10:1206-1213.
- 18 49 Mednick SC, Cai DJ, Kanady J, Drummond SPA: Comparing the benefits of
19 caffeine, naps and placebo on verbal, motor and perceptual memory. *Behav Brain*
20 *Res* 2008;193:79-86.
- 21 50 Morin A, Doyon J, Dostie V, Barakat M, Tahar AH, Korman M, Benali H,
22 Karni A, Ungerleider LG, Carrier J: Motor sequence learning increases sleep
23 spindles and fast frequencies in post-training sleep. *Sleep* 2008;31:1149-1156.
- 24 51 Nishida M, Walker MP: Daytime naps, motor memory consolidation and
25 regionally specific sleep spindles. *PLoS ONE* 2007;2:e341.
- 26 52 Rasch B, Büchel C, Gais S, Born J: Odor cues during slow-wave sleep prompt
27 declarative memory consolidation. *Science* 2007;315:1426-1429.
- 28 53 Sheth BR, Janvelyan D, Khan M: Practice makes imperfect: Restorative effects
29 of sleep on motor learning. *PLoS ONE* 2008;3:e3190.
- 30 54 Walker MP, Brakefield T, Hobson JA, Stickgold R: Dissociable stages of
31 human memory consolidation and reconsolidation. *Nature* 2003;425:616-620.
- 32 55 Walker MP, Brakefield T, Seidman J, Morgan A, Hobson JA, Stickgold R:
33 Sleep and the time course of motor skill learning. *Learn Mem* 2003;10:275-284.
- 34 56 Silverman I, Phillips K: Effects of estrogen changes during the menstrual cycle
35 on spatial performance. *Ethology and Sociobiology* 1993;14:257-269.
- 36 57 Szekely C, Hampson E, Carey DP, Goodale MA: Oral contraceptive use affects
37 manual praxis but not simple visually guided movements. *Developmental*
38 *Neuropsychology* 1998;14:399-420.

- 1 58 Silber M, Almkvist O, Larsson B, Stock S, Uvn s-Moberg K: The effect of
2 oral contraceptive pills on levels of oxytocin in plasma and on cognitive functions.
3 Contraception 1987;36:641-650.
- 4 59 Beck KD, McLaughlin J, Bergen MT, Cominski TP, Moldow RL, Servatius
5 RJ: Facilitated acquisition of the classically conditioned eyeblink response in women
6 taking oral contraceptives. Behavioural Pharmacology 2008;19.
- 7 60 Wilhelm I, Metzkw-M  sz  ros M, Knapp S, Born J: Sleep-dependent
8 consolidation of procedural motor memories in children and adults: the pre-sleep
9 level of performance matters. Developmental Science 2012;15:506-515.
- 10 61 Galea LAM, Uban KA, Epp JR, Brummelte S, Barha CK, Wilson WL,
11 Lieblich SE, Pawluski JL: Endocrine regulation of cognition and neuroplasticity:
12 Our pursuit to unveil the complex interaction between hormones, the brain, and
13 behaviour. Canadian Journal of Experimental Psychology/Revue canadienne de
14 psychologie exp  rimentale 2008;62:247-260.
- 15 62 Spencer JL, Waters EM, Romeo R, Wood GE, Milner TA, McEwen B:
16 Uncovering the mechanisms of estrogen effects on hippocampal function. Frontiers
17 in Neuroendocrinology 2008;29:219-237.
- 18 63 McEwen B: Estrogen Actions Throughout the Brain. Recent Prog Horm Res
19 2002;57:357-384.
- 20 64 Romeo RD, Waters EM, McEwen BS: Steroid-induced hippocampal synaptic
21 plasticity: sex differences and similarities. Neuron Glia Biology 2004;1:219-229.
- 22 65 Jazin E, Cahill L: Sex differences in molecular neuroscience: from fruit flies to
23 humans. Nat Rev Neurosci 2010;11:9-17.
- 24 66 Sandstrom NJ, Kaufman J, Huettel A: Males and females use different distal
25 cues in a virtual environment navigation task. Cognitive Brain Research 1998;6:351-
26 360.
- 27 67 Lewin C, Wolgers G, Herlitz A: Sex differences favoring women in verbal but
28 not in visuospatial episodic memory. Neuropsychology 2001;15:165-173.
- 29 68 Postma A, Winkel J, Tuiten A, van Honk J: Sex differences and menstrual
30 cycle effects in human spatial memory. Psychoneuroendocrinology 1999;24:175-192.
- 31 69 Weiss EM, Kemmler G, Deisenhammer EA, Fleischhacker WW, Delazer M:
32 Sex differences in cognitive functions. Pers Indiv Differ 2003;35:863-875.
- 33 70 Cahill L: Why sex matters for neuroscience. Nat Rev Neurosci 2006;7:477-484.
- 34 71 Hampson E: Variations in sex-related cognitive abilities across the menstrual
35 cycle. Brain Cogn 1990;14:26-43.
- 36 72 Hampson E: Reciprocal effects of hormonal fluctuations on human motor and
37 perceptual-spatial skills. Behav Neurosci 1988;102:456-459.
38

1

2





Figure 1: Study design: All participants underwent 6 study days: nap with learning (L-Nap), wake with learning (L-Wake), nap without learning (C-Nap); each of the three conditions once in the active OC week (second week of the three pill weeks) and once in the OC free week. The order of all 6 conditions was balanced across participants.

1



2

3

Figure 2: Change in declarative (absolute change in number of words, with SEM, left) and

4

motor (relative change in correctly tapped sequences during 30 sec trial, with SEM, right)

5

performance from the learning phase (13:00 h) to the retest in the afternoon (16:30 h),

6

separated in the groups with (L-Nap) and without (L-Wake) a nap between learning and retest.

7

There was no significant difference between the conditions.

8

Table 1: Absolute task performance at the end of training and at retest after either nap (L-NAP) or wake (L-Wake) during the active OC and OC free week (mean with SD). For both tasks a significant increase in performance was seen after the offline period regardless of nap/wake condition or OC phase. All tests were significant after correction for multiple comparisons for each task ($p < 0.0125$).

		Active OC week		OC free week	
		L-NAP	L-Wake	L-NAP	L-Wake
Word Pairs	end-training	27.2 ± 7.9	27.0 ± 7.5	27.6 ± 7.7	26.7 ± 5.5
	retest	35.5 ± 4.8	34.2 ± 3.6	35.3 ± 3.6	35.3 ± 3.3
	statistics	$T_{14}=6.6; P<.001$	$T_{14}=5.7; P<.001$	$T_{14}=5.8; P<.001$	$T_{14}=7.6; P<.001$
Tapping	end-training	18.2 ± 3.9	18.7 ± 3.9	18.9 ± 2.9	18.3 ± 3.5
	retest	21.2 ± 4.6	21.6 ± 5.0	20.9 ± 3.7	20.4 ± 3.4
	statistics	$T_{14}=7.3; P<.001$	$T_{14}=5.7; P<.001$	$T_{14}=3.5; P=.003$	$T_{14}=3.1; P=.007$

Table 2: Sleep stage duration (minutes with SD) and power in the EEG frequency bands (μV^2 with SD) of the nap with (L-NAP) and without (C-NAP) previous learning session. Data are reported as obtained during the OC week and during the OC free week. There was no significant difference between the two conditions and the two weeks.

	Active OC week		OC free week		Statistics
	L-NAP	C-NAP	L-NAP	C-NAP	Repeated Measures MANOVA
(min)					
S1	11.3 ± 7.5	14.3 ± 10.9	11.1 ± 9.2	11.5 ± 8.0	Nap: F _{5,10} =.720; p>.6 Week: F _{5,10} =.535; p>.7 Nap* Week: F _{5,10} =1.264; p>.3
S2	31.2 ± 18.3	26.5 ± 13.1	28.3 ± 13.6	30.3 ± 9.2	
SWS	20.0 ± 10.4	20.2 ± 12.4	23.1 ± 20.3	19.3 ± 13.1	
REM	2.6 ± 5.8	1.8 ± 3.6	2.9 ± 4.9	5.3 ± 6.8	
TST	66.1 ± 14.6	62.9 ± 14.3	65.6 ± 24.0	66.5 ± 12.7	
(μV ²)					
Delta	550 ± 208	687 ± 452	605 ± 490	581 ± 288	Nap: F _{5,10} =1.255; p>.3 Week: F _{5,10} =1.141; p=.4 Nap* Week: F _{5,10} =.537; p>.7
Theta	87 ± 35	102 ± 54	79 ± 24	90 ± 42	
Alpha	52 ± 24	63 ± 45	49 ± 26	51 ± 23	
Sigma	23 ± 13	27 ±17	20 ± 8	23 ± 10	
Beta	8 ± 3	9 ± 7	8 ± 5	8 ± 4	

Table 3: Endogenous hormone values (mean with SD in pg/ml) for all conditions: nap with learning (L-NAP), wake with learning (L-WAKE), nap without learning (C-NAP).

	17-Beta Estrogen			Progesterone		
	L-Nap	C-Nap	L-Wake	L-Nap	C-Nap	L-Wake
OC week	12.0 ± 8.2	9.5 ± 4.4	8.9 ± 4.2	.30 ± .16	.25 ± .18	.28 ± .18
OC free week	36.0 ± 49.0	48.6 ± 41.9	25.8 ± 23.4	.32 ± .19	.28 ± .20	.30 ± .16
statistics	T ₁₄ =1.9;	T ₁₄ =3.5;	T ₁₄ =2.8;	T ₁₄ =.59;	T ₁₄ =1.2;	T ₁₄ =.72;
	P=.08	P<.005	P<.02	P>.5	P>.25	P>.45